



MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

MATH 122

Blake Farman ¹

¹University of South Carolina, Columbia, SC USA

Calculus for Business Administration and Social
Sciences



OUTLINE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

1 6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS



OUTLINE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

1 6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

2 6.6: INTEGRATION BY SUBSTITUTION

- Examples



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^3 2x dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int_1^3 2x dx.$$

$$\int_1^3 2x dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int_1^3 2x dx.$$

$$\int_1^3 2x dx = 2 \int_1^3 x dx$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^3 2x dx.$$

$$\begin{aligned}\int_1^3 2x dx &= 2 \int_1^3 x dx \\ &= 2 \left[\frac{1}{2} x^2 \right]_1^3\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_1^3 2x dx.$$

$$\begin{aligned}\int_1^3 2x dx &= 2 \int_1^3 x dx \\ &= 2 \left[\frac{1}{2} x^2 \right]_1^3 \\ &= x^2 \Big|_1^3\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_1^3 2x dx.$$

$$\begin{aligned}\int_1^3 2x dx &= 2 \int_1^3 x dx \\ &= 2 \left[\frac{1}{2} x^2 \right]_1^3 \\ &= x^2 \Big|_1^3 \\ &= 3^2 - 1^2\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_1^3 2x \, dx.$$

$$\begin{aligned}\int_1^3 2x \, dx &= 2 \int_1^3 x \, dx \\ &= 2 \left[\frac{1}{2} x^2 \right]_1^3 \\ &= x^2 \Big|_1^3 \\ &= 3^2 - 1^2 \\ &= 9 - 1\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_1^3 2x \, dx.$$

$$\begin{aligned}\int_1^3 2x \, dx &= 2 \int_1^3 x \, dx \\ &= 2 \left[\frac{1}{2} x^2 \right]_1^3 \\ &= x^2 \Big|_1^3 \\ &= 3^2 - 1^2 \\ &= 9 - 1 \\ &= 8.\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\int_0^2 6x^2 dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\int_0^2 6x^2 dx = 6 \int_0^2 x^2 dx$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\begin{aligned}\int_0^2 6x^2 dx &= 6 \int_0^2 x^2 dx \\ &= \left. \frac{6}{3} x^3 \right|_0^2\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\begin{aligned}\int_0^2 6x^2 dx &= 6 \int_0^2 x^2 dx \\ &= \frac{6}{3} x^3 \Big|_0^2 \\ &= 2(2^3 - 0^3)\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\begin{aligned}\int_0^2 6x^2 dx &= 6 \int_0^2 x^2 dx \\ &= \frac{6}{3} x^3 \Big|_0^2 \\ &= 2(2^3 - 0^3) \\ &= 2(8)\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^2 6x^2 dx.$$

$$\begin{aligned}\int_0^2 6x^2 dx &= 6 \int_0^2 x^2 dx \\ &= \frac{6}{3} x^3 \Big|_0^2 \\ &= 2(2^3 - 0^3) \\ &= 2(8) \\ &= 16.\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^2 t^3 dt.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION
EXAMPLES

Compute

$$\int_0^2 t^3 dt.$$

$$\int_0^2 t^3 dt =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION
EXAMPLES

Compute

$$\int_0^2 t^3 dt.$$

$$\int_0^2 t^3 dt = \left. \frac{1}{4}t^4 \right|_0^2$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION
EXAMPLES

Compute

$$\int_0^2 t^3 dt.$$

$$\begin{aligned}\int_0^2 t^3 dt &= \left. \frac{1}{4} t^4 \right|_0^2 \\ &= \frac{1}{4} (16 - 0)\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^2 t^3 dt.$$

$$\begin{aligned}\int_0^2 t^3 dt &= \left. \frac{1}{4} t^4 \right|_0^2 \\ &= \frac{1}{4}(16 - 0) \\ &= 4.\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\int_1^2 8x + 5 dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\int_1^2 8x + 5 dx = 8 \int_1^2 x dx + 5 \int_1^2 dx$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\begin{aligned} \int_1^2 8x + 5 dx &= 8 \int_1^2 x dx + 5 \int_1^2 dx \\ &= \left. \frac{8}{2}x^2 \right|_1^2 + \left. 5x \right|_1^2 \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\begin{aligned}\int_1^2 8x + 5 dx &= 8 \int_1^2 x dx + 5 \int_1^2 dx \\ &= \left. \frac{8}{2}x^2 \right|_1^2 + \left. 5x \right|_1^2 \\ &= 4(4 - 1) + 5(2 - 1)\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\begin{aligned}\int_1^2 8x + 5 dx &= 8 \int_1^2 x dx + 5 \int_1^2 dx \\ &= \left. \frac{8}{2}x^2 \right|_1^2 + \left. 5x \right|_1^2 \\ &= 4(4 - 1) + 5(2 - 1) \\ &= 12 + 5\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int_1^2 8x + 5 dx.$$

$$\begin{aligned} \int_1^2 8x + 5 dx &= 8 \int_1^2 x dx + 5 \int_1^2 dx \\ &= \left. \frac{8}{2}x^2 + 5x \right|_1^2 \\ &= 4(4 - 1) + 5(2 - 1) \\ &= 12 + 5 \\ &= 17. \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$

$$\int_0^1 8e^{2t} dt =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$

$$\int_0^1 8e^{2t} dt = 8 \int_0^1 e^{2t} dt$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$

$$\begin{aligned}\int_0^1 8e^{2t} dt &= 8 \int_0^1 e^{2t} dt \\ &= \frac{8}{2} e^{2t} \Big|_0^1\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$

$$\begin{aligned}\int_0^1 8e^{2t} dt &= 8 \int_0^1 e^{2t} dt \\ &= \frac{8}{2} e^{2t} \Big|_0^1 \\ &= 4(e^2 - e^0)\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int_0^1 8e^{2t} dt.$$

$$\begin{aligned}\int_0^1 8e^{2t} dt &= 8 \int_0^1 e^{2t} dt \\ &= \frac{8}{2} e^{2t} \Big|_0^1 \\ &= 4(e^2 - e^0) \\ &= 4(e^2 - 1).\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} = e^{x^2} \cdot 2x$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} = e^{x^2} \cdot 2x = e^u \cdot \frac{du}{dx}.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} = e^{x^2} \cdot 2x = e^u \cdot \frac{du}{dx}.$$

This looks exactly like the result of applying the Chain Rule to the composition $e^{u(x)}$!



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} = e^{x^2} \cdot 2x = e^u \cdot \frac{du}{dx}.$$

This looks exactly like the result of applying the Chain Rule to the composition $e^{u(x)}$! This tells us that

$$\int 2xe^{x^2} dx = e^{x^2} + c.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

To compute

$$\int 2xe^{x^2} dx$$

we must find an antiderivative. We note that $\frac{d}{dx}x^2 = 2x$. If we let $u = x^2$, then

$$2xe^{x^2} = e^{x^2} \cdot 2x = e^u \cdot \frac{du}{dx}.$$

This looks exactly like the result of applying the Chain Rule to the composition $e^{u(x)}$! This tells us that

$$\int 2xe^{x^2} dx = e^{x^2} + c.$$

The method of Integration by Substitution is intended to integrate a product of functions that appears to have come from an application of the chain rule.



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function.



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u .



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral.



THE METHOD

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral. In this example,



THE METHOD

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral. In this example,

$$\int 2xe^{x^2} dx =$$



THE METHOD

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral. In this example,

$$\int 2xe^{x^2} dx = \int e^u du$$



THE METHOD

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral. In this example,

$$\begin{aligned} \int 2xe^{x^2} dx &= \int e^u du \\ &= e^u + c \end{aligned}$$



THE METHOD

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

- 1 Identify a piece of the integrand that looks like it could be the derivative of another piece of the integrand, and we let u be that function. Here, $u = 2x$.
- 2 We formally treat dx and du like variables to change variables from x to u . Here, we multiply both sides of $2xe^{x^2} = e^u \frac{du}{dx}$ by dx to obtain

$$2xe^{x^2} dx = e^u du$$

- 3 Finally, since these expressions are equal, we can evaluate the simpler integral. In this example,

$$\begin{aligned}\int 2xe^{x^2} dx &= \int e^u du \\ &= e^u + c \\ &= e^{x^2} + c.\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

- 1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

- 1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.
- 2 We have



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

2 We have

$$\frac{du}{dx} =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

① We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

② We have

$$\frac{du}{dx} = 2x$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

- 1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.
- 2 We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

2 We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$

3 Compute the simplified integral:



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

① We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

② We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$

③ Compute the simplified integral:

$$\int (x^2 + 1)^5 2x \, dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

2 We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$

3 Compute the simplified integral:

$$\int (x^2 + 1)^5 2x \, dx = \int u^5 \, du$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

2 We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$

3 Compute the simplified integral:

$$\begin{aligned} \int (x^2 + 1)^5 2x \, dx &= \int u^5 \, du \\ &= \frac{1}{6} u^6 + c \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int (x^2 + 1)^5 \cdot 2x \, dx.$$

1 We note that $2x = \frac{d}{dx}(x^2 + 1)$, so we let $u = x^2 + 1$.

2 We have

$$\frac{du}{dx} = 2x \Rightarrow du = 2x \, dx.$$

3 Compute the simplified integral:

$$\begin{aligned} \int (x^2 + 1)^5 2x \, dx &= \int u^5 \, du \\ &= \frac{1}{6} u^6 + c \\ &= \frac{1}{6} (x^2 + 1)^6 + c. \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$

Therefore

$$\int \frac{2x}{x^2 + 4} dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$

Therefore

$$\int \frac{2x}{x^2 + 4} dx = \int \frac{1}{u} du$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$

Therefore

$$\begin{aligned} \int \frac{2x}{x^2 + 4} dx &= \int \frac{1}{u} du \\ &= \ln |u| + c \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$

Therefore

$$\begin{aligned} \int \frac{2x}{x^2 + 4} dx &= \int \frac{1}{u} du \\ &= \ln |u| + c \\ &= \ln |x^2 + 4| + c \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{2x}{x^2 + 4} dx.$$

Let $u = x^2 + 4$ so

$$\frac{du}{dx} = 2x \Rightarrow du = 2x dx.$$

Therefore

$$\begin{aligned} \int \frac{2x}{x^2 + 4} dx &= \int \frac{1}{u} du \\ &= \ln |u| + c \\ &= \ln |x^2 + 4| + c \\ &= \ln(x^2 + 4) + c. \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore

$$\int te^{t^2+1} dt =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore

$$\int te^{t^2+1} dt = \int e^u \frac{du}{2}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore

$$\begin{aligned}\int te^{t^2+1} dt &= \int e^u \frac{du}{2} \\ &= \frac{1}{2} \int e^u du\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore

$$\begin{aligned}\int te^{t^2+1} dt &= \int e^u \frac{du}{2} \\ &= \frac{1}{2} \int e^u du \\ &= \frac{1}{2} e^u + c\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int te^{t^2+1} dt.$$

Let $u = t^2 + 1$ so $du = 2t dt$. Since we want to replace $t dt$, we divide both sides by 2 to get $du/2 = t dt$. Therefore

$$\begin{aligned}\int te^{t^2+1} dt &= \int e^u \frac{du}{2} \\ &= \frac{1}{2} \int e^u du \\ &= \frac{1}{2} e^u + c \\ &= \frac{1}{2} e^{t^2+1} + c.\end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.
Therefore



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.
Therefore

$$\int x^3 \sqrt{x^4 + 5} dx =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION

EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.
Therefore

$$\int x^3 \sqrt{x^4 + 5} dx = \int \sqrt{u} \frac{du}{4}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.
Therefore

$$\begin{aligned} \int x^3 \sqrt{x^4 + 5} dx &= \int \sqrt{u} \frac{du}{4} \\ &= \frac{1}{4} \int u^{1/2} du \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4x^3 dx$. Hence $du/4 = x^3 dx$.
Therefore

$$\begin{aligned} \int x^3 \sqrt{x^4 + 5} dx &= \int \sqrt{u} \frac{du}{4} \\ &= \frac{1}{4} \int u^{1/2} du \\ &= \frac{1}{4} \left(\frac{2}{3} u^{3/2} \right) + c \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int x^3 \sqrt{x^4 + 5} dx.$$

Let $u = x^4 + 5$ so $du = 4^3 x dx$. Hence $du/4 = x^3 dx$.
Therefore

$$\begin{aligned} \int x^3 \sqrt{x^4 + 5} dx &= \int \sqrt{u} \frac{du}{4} \\ &= \frac{1}{4} \int u^{\frac{1}{2}} du \\ &= \frac{1}{4} \left(\frac{2}{3} u^{\frac{3}{2}} \right) + c \\ &= \frac{1}{6} (x^4 + 5)^{\frac{3}{2}} + c. \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDA-
MENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITU-
TION

EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION

EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore

$$\int \frac{t^2}{1+t^3} dt =$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore

$$\int \frac{t^2}{1+t^3} dt = \int \frac{1}{u} \frac{du}{3}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING
THE FUNDAMENTAL
THEOREM TO
COMPUTE
DEFINITE
INTEGRALS

6.6:
INTEGRATION
BY SUBSTITUTION
EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore

$$\begin{aligned} \int \frac{t^2}{1+t^3} dt &= \int \frac{1}{u} \frac{du}{3} \\ &= \frac{1}{3} \int \frac{du}{u} \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore

$$\begin{aligned} \int \frac{t^2}{1+t^3} dt &= \int \frac{1}{u} \frac{du}{3} \\ &= \frac{1}{3} \int \frac{du}{u} \\ &= \ln |u| + c \end{aligned}$$



EXAMPLE

MATH 122

FARMAN

6.3: USING THE FUNDAMENTAL THEOREM TO COMPUTE DEFINITE INTEGRALS

6.6: INTEGRATION BY SUBSTITUTION
EXAMPLES

Compute

$$\int \frac{t^2}{1+t^3} dt.$$

Let $u = 1 + t^3$ so $du = 3t^2 dt$. Hence $du/3 = t^2 dt$.
Therefore

$$\begin{aligned} \int \frac{t^2}{1+t^3} dt &= \int \frac{1}{u} \frac{du}{3} \\ &= \frac{1}{3} \int \frac{du}{u} \\ &= \ln |u| + c \\ &= \ln |1+t^3| + c. \end{aligned}$$